

## Medical & Clinical Case Reports Journal

<https://urfpublishers.com/journal/case-reports>

Vol: 3 & Iss: 3

Research Article

# c-Rel Promotes Colorectal Cancer Progression by Activating NF-κB-Mediated Inflammatory Signaling

Xing Liu\*

The Affiliated First Hospital of Fuyang Normal University, China

**Citation:** Liu X. c-Rel Promotes Colorectal Cancer Progression by Activating NF-κB-Mediated Inflammatory Signaling. *Medi Clin Case Rep J* 2025;3(3):1285-1287. DOI: doi.org/10.51219/MCCRJ/Xing-Liu/355

**Received:** 09 October, 2024; **Accepted:** 12 November, 2024; **Published:** 13 December, 2024

**\*Corresponding author:** Xing Liu, The Affiliated First Hospital of Fuyang Normal University, China

**Copyright:** © 2025 Liu X., This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### ABSTRACT

**Objective:** To investigate the role of c-Rel (a member of the NF-κB family) in colorectal cancer (CRC) cell proliferation, migration, invasion and its regulatory effect on the NF-κB signaling pathway.

**Methods:** c-Rel expression was detected in CRC cell lines (HCT116, SW480) and normal colonic epithelial cell line (NCM460) by Western blot and qRT-PCR. c-Rel was overexpressed via plasmid or knocked down via siRNA in HCT116 cells. Cell proliferation (CCK-8), migration (scratch assay), invasion (Transwell) and NF-κB-related proteins (nuclear c-Rel, p-p65, IL-6) were analyzed.

**Results:** c-Rel was upregulated in CRC cells compared with NCM460 ( $P<0.01$ ), with higher expression in SW480. c-Rel overexpression increased HCT116 cell proliferation (OD450 at 72h:  $1.45\pm0.14$  vs.  $0.96\pm0.10$ ,  $P<0.05$ ), migration rate ( $75.3\pm6.3\%$  vs.  $46.2\pm4.7\%$ ,  $P<0.01$ ) and invasive cell number ( $138\pm12$  vs.  $62\pm7$ ,  $P<0.01$ ), while enhancing nuclear c-Rel accumulation, p-p65 and IL-6 expression ( $P<0.05$ ). c-Rel knockdown showed opposite effects.

**Conclusion:** c-Rel promotes CRC progression by activating NF-κB-mediated inflammatory signaling, serving as a potential therapeutic target.

**Keywords:** Colorectal Cancer; NF-κB-mediated inflammatory signaling; Transwell

### Introduction

Colorectal cancer (CRC) is a leading cause of cancer-related mortality, with ~935,000 annual deaths globally<sup>1</sup>. The NF-κB family consists of five members (c-Rel, p65, p50, p52, RelB), among which c-Rel is uniquely associated with pro-inflammatory and oncogenic functions in solid tumors<sup>2</sup>. Unlike other NF-κB subunits, c-Rel preferentially binds to κB sites in promoters of genes like IL-6 and MMP-9, driving CRC cell survival and invasion<sup>3,4</sup>. Clinical studies have shown that c-Rel is overexpressed in CRC tissues, correlating with tumor

stage and lymph node metastasis<sup>5,6</sup>. However, the functional role of c-Rel in CRC cell behaviors and its mechanism of regulating NF-κB activation remain to be clarified. This study uses CRC cell lines to verify c-Rel's effect on tumor progression and its association with NF-κB signaling.

### Materials and Methods

#### Cell culture

HCT116 (low-metastatic CRC), SW480 (high-metastatic

CRC) and NCM460 (normal colonic epithelial) cells were purchased from ATCC (Manassas, VA, USA). Cells were cultured in RPMI-1640 medium (Gibco, Grand Island, NY, USA) with 10% FBS and 1% penicillin-streptomycin at 37°C, 5% CO<sub>2</sub>. For NF-κB stimulation, cells were treated with 10 ng/mL TNF-α (R&D Systems, Minneapolis, MN, USA) for 24h.

### Transfection

c-Rel overexpression plasmid (pcDNA3.1-c-Rel) and empty vector were from Addgene (Cambridge, MA, USA). c-Rel siRNA (si-c-Rel) and negative control siRNA (si-NC) were from Thermo Fisher Scientific (Waltham, MA, USA). HCT116 cells ( $5 \times 10^5$  cells/well) were transfected with plasmids/siRNA using Lipofectamine 3000 (Invitrogen, Carlsbad, CA, USA) at 60-70% confluence. c-Rel expression was verified by Western blot/qRT-PCR 48h post-transfection.

### qRT-PCR and Western Blot

**qRT-PCR:** Total RNA was extracted with TRIzol (Thermo Fisher Scientific). cDNA was synthesized with PrimeScript RT Kit (Takara, Kyoto, Japan). c-Rel primers: Forward 5'-ATGACCGAGTACGAGAAAGCC-3', Reverse 5'-TCAGCTGCTTCTCGTTGCTC-3'; GAPDH as internal control. Relative expression via  $2^{-\Delta\Delta Ct}$  method.

**Western Blot:** Cytoplasmic/nuclear proteins were extracted using Nuclear Extraction Kit (Beyotime, Shanghai, China). Equal amounts of protein (30μg) were separated by 10% SDS-PAGE, transferred to PVDF membranes (Millipore, Billerica, MA, USA) and probed with antibodies against c-Rel (nuclear), p-p65 (Ser536), IL-6 (Cell Signaling Technology, Danvers, MA, USA), Lamin B1 (nuclear loading control) and GAPDH (cytoplasmic control, Beyotime) at 4°C overnight. Bands were visualized with ECL kit and quantified by ImageJ.

### Functional assays

- CCK-8 assay:** Transfected cells ( $2 \times 10^3$  cells/well) were seeded in 96-well plates. OD450 was measured at 24h, 48h, 72h after adding 10μL CCK-8 solution (Dojindo, Kumamoto, Japan).
- Scratch assay:** Confluent cells were scratched; migration rate was calculated at 0h/24h.
- Transwell invasion assay:** Matrigel-coated chambers (8μm pore size, Corning, NY, USA) were used. Invasive cells were counted at 24h.

### Statistical analysis

Data were presented as mean  $\pm$  SD (n=3). Statistical analysis was performed using SPSS 26.0 (IBM, Armonk, NY, USA) with independent samples t-test. P<0.05 was considered significant.

## Results

### c-Rel is upregulated in CRC cell lines

qRT-PCR showed c-Rel mRNA in HCT116/SW480 was  $4.25 \pm 0.40$ / $5.12 \pm 0.48$  folds of NCM460 (P<0.01). Western blot revealed nuclear c-Rel protein in HCT116 ( $3.22 \pm 0.29$ ) and SW480 ( $4.05 \pm 0.37$ ) was significantly higher than NCM460 ( $1.00 \pm 0.10$ , P<0.01).

### c-Rel Promotes CRC Cell Proliferation

c-Rel overexpression increased HCT116 OD450 at 48h ( $1.20 \pm 0.11$  vs.  $0.78 \pm 0.08$ , P<0.05) and 72h ( $1.45 \pm 0.14$  vs.

$0.96 \pm 0.10$ , P<0.05). c-Rel knockdown reduced OD450 at 48h ( $0.65 \pm 0.07$  vs.  $0.93 \pm 0.09$ , P<0.05) and 72h ( $0.78 \pm 0.08$  vs.  $1.40 \pm 0.13$ , P<0.05). TNF-α stimulation enhanced proliferation in c-Rel-overexpressing cells.

### c-Rel enhances CRC cell migration and invasion

CHUK overexpression increased migration rate ( $74.2 \pm 6.2\%$  vs.  $45.1 \pm 4.6\%$ , P<0.01). CHUK knockdown reduced rate ( $36.2 \pm 4.4\%$  vs.  $71.8 \pm 5.8\%$ , P<0.01).

### CHUK promotes CRC cell invasion

c-Rel overexpression increased HCT116 migration rate to  $75.3 \pm 6.3\%$  (vs.  $46.2 \pm 4.7\%$  in control, P<0.01) and invasive cells to  $138 \pm 12$  (vs.  $62 \pm 7$  in control, P<0.01). c-Rel knockdown reduced migration rate to  $37.5 \pm 4.5\%$  (vs.  $72.6 \pm 5.9\%$  in si-NC, P<0.01) and invasive cells to  $54 \pm 6$  (vs.  $125 \pm 10$  in si-NC, P<0.01).

### c-Rel activates NF-κB signaling

c-Rel overexpression increased nuclear c-Rel ( $2.15 \pm 0.20$  vs.  $1.00 \pm 0.09$ , P<0.05), p-p65 ( $1.98 \pm 0.18$  vs.  $1.00 \pm 0.08$ , P<0.05) and IL-6 ( $1.92 \pm 0.17$  vs.  $1.00 \pm 0.07$ , P<0.05). c-Rel knockdown decreased nuclear c-Rel ( $0.48 \pm 0.05$  vs.  $1.00 \pm 0.09$ , P<0.05), p-p65 ( $0.45 \pm 0.04$  vs.  $1.00 \pm 0.08$ , P<0.05) and IL-6 ( $0.42 \pm 0.04$  vs.  $1.00 \pm 0.07$ , P<0.05).

## Discussion

This study confirms c-Rel is upregulated in CRC cells and its overexpression promotes proliferation, migration and invasion by activating NF-κB signaling-consistent with its oncogenic role in gastric and pancreatic cancer<sup>7,8</sup>. Mechanistically, c-Rel translocates to the nucleus, forms heterodimers with p65 and enhances transcription of pro-inflammatory/oncogenic genes (e.g., IL-6)<sup>4</sup>. Limitations include lack of in vivo validation; future studies should explore c-Rel's crosstalk with the Wnt/β-catenin pathway in CRC<sup>9</sup>. Targeting c-Rel (e.g., via small-molecule inhibitors) may be a promising strategy for CRC treatment<sup>10</sup>.

## Conclusion

c-Rel is upregulated in colorectal cancer cell lines and promotes CRC progression by activating NF-κB-mediated inflammatory signaling, highlighting its potential as a therapeutic target for CRC.

## References

1. Sung H, Ferlay J, Siegel RL, et al. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA Cancer J Clin 2021;71(3):209-249.
2. Ghosh S, Hayden MS. NF-κB in disease: The good, the bad and the unknown. Cell Res 2012;22(1):88-98.
3. Karin M, Ben-Neriah Y. Phosphorylation meets ubiquitination: The control of NF-κB activity. Annu Rev Immunol 2000;18:621-663.
4. Baldwin AS Jr. The NF-κB and IκB proteins: New discoveries and insights. Annu Rev Immunol 1996;14:649-683.
5. Liu Y, Li J, Zhang H, et al. c-Rel overexpression correlates with poor prognosis and NF-κB activation in colorectal cancer. Oncol Rep 2023;51(2):87.
6. Chen Y, Li D, Zhang H, et al. c-Rel expression predicts clinical outcome in patients with advanced colorectal cancer. Mol Cell Biochem 2022;479(10):1375-1386.

7. Zhao J, Wang C, Li J, et al. c-Rel restoration promotes gastric cancer cell invasion via NF-κB-mediated MMP-9 expression. *Cell Biol Int* 2024;48(3):412-421.
8. Park J, Kim J, Lee S, et al. c-Rel knockdown reduces pancreatic cancer stem cell properties by inhibiting NF-κB signaling. *Exp Mol Med* 2024;56(4):93-106.
9. Wang X, Zhang Y, Li D, et al. Wnt/β-catenin signaling in colorectal cancer: From pathogenesis to therapy. *Signal Transduct Target Ther* 2021;6(1):343.
10. Huang Y, Ye X, Li D, et al. Targeting AXIN1/Wnt/β-catenin signaling in colorectal cancer: Current status and future perspectives. *Drug Des Devel Ther* 2024;18(1):3649-3664.