

## Medical & Clinical Case Reports Journal

<https://urfpublishers.com/journal/case-reports>

Vol: 3 & Iss: 3

Research Article

# NFKBIA Inhibits Colorectal Cancer Progression via Suppressing the NF-κB Signaling Pathway

Xing Liu\*

The Affiliated First Hospital of Fuyang Normal University, China

**Citation:** Liu X. NFKBIA Inhibits Colorectal Cancer Progression via Suppressing the NF-κB Signaling Pathway. *Medi Clin Case Rep J* 2025;3(3):1333-1335. DOI: doi.org/10.51219/MCCRJ/Xing-Liu/371

**Received:** 03 February, 2025; **Accepted:** 05 March, 2025; **Published:** 07 April, 2025

**\*Corresponding author:** Xing Liu, The Affiliated First Hospital of Fuyang Normal University, China

**Copyright:** © 2025 Liu X., This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### ABSTRACT

**Objective:** To investigate the role of NFKBIA (nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha, also known as IκBα) in colorectal cancer (CRC) cell proliferation, migration, invasion and its regulation of the NF-κB signaling pathway.

**Methods:** NFKBIA expression in CRC cell lines (HCT116, SW480) and normal colonic epithelial cell line (NCM460) was detected by Western blot and qRT-PCR. NFKBIA was overexpressed via plasmid or knocked down via siRNA in HCT116 cells. Cell proliferation (CCK-8), migration (scratch assay), invasion (Transwell) and NF-κB-related proteins (p-p65, IκBα, TNF-α) were analyzed.

**Results:** NFKBIA was downregulated in CRC cells ( $P<0.01$ ). NFKBIA overexpression reduced proliferation (OD<sub>450</sub> at 72h:  $0.66\pm0.06$  vs.  $1.30\pm0.12$ ,  $P<0.05$ ), migration (24h rate:  $29.8\pm3.7\%$  vs.  $68.5\pm5.6\%$ ,  $P<0.01$ ), invasion (cell number:  $41\pm5$  vs.  $124\pm10$ ,  $P<0.01$ ) and downregulated p-p65, TNF-α ( $P<0.05$ ). NFKBIA knockdown showed opposite effects.

**Conclusion:** NFKBIA suppresses CRC progression via inhibiting NF-κB signaling, serving as a potential therapeutic target.

**Keywords:** Colorectal Cancer; Cell Proliferation; Transwell; Nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha

### Introduction

Colorectal cancer (CRC) causes ~935,000 annual deaths globally, with constitutively activated NF-κB signaling being a key driver of its inflammatory progression<sup>1</sup>. NFKBIA (IκBα) is the major endogenous inhibitor of NF-κB: it binds to cytoplasmic p65/p50 complexes to prevent nuclear translocation, thereby blocking NF-κB-mediated oncogenic gene expression<sup>2,3</sup>. NFKBIA is frequently downregulated in gastric, pancreatic and CRC, correlating with high NF-κB activity and poor prognosis<sup>4,5</sup>.

However, NFKBIA's functional role in regulating CRC cell behaviors and its impact on NF-κB suppression remain to be clarified. This study explores NFKBIA's effect on CRC cells and its association with the NF-κB signaling axis.

### Materials and Methods

#### Cell culture

HCT116, SW480 (CRC cell lines) and NCM460 (normal colonic epithelial cell line) were purchased from ATCC

(Manassas, VA, USA). Cells were cultured in RPMI-1640 medium (Gibco, Grand Island, NY, USA) supplemented with 10% fetal bovine serum (FBS) and 1% penicillin-streptomycin at 37°C in a 5% CO<sub>2</sub> humidified incubator. For NF-κB stimulation, cells were treated with 10 ng/mL TNF-α (R&D Systems, Minneapolis, MN, USA) for 24h.

### Transfection

NFKBIA overexpression plasmid (pcDNA3.1-NFKBIA) and empty vector were obtained from Addgene (Cambridge, MA, USA). NFKBIA siRNA (si-NFKBIA) and negative control siRNA (si-NC) were purchased from Thermo Fisher Scientific (Waltham, MA, USA). HCT116 cells (5×10<sup>5</sup> cells/well) were seeded in 6-well plates and transfected with plasmids or siRNA using Lipofectamine 3000 (Invitrogen, Carlsbad, CA, USA) at 60-70% confluence. NFKBIA expression was verified by Western blot and qRT-PCR 48h post-transfection.

### qRT-PCR and western blot

**qRT-PCR:** Total RNA was extracted with TRIzol reagent (Thermo Fisher Scientific). cDNA was synthesized using PrimeScript RT Kit (Takara, Kyoto, Japan). NFKBIA primers: Forward 5'-GCTGCTGCTGCTGTTCTGA-3', Reverse 5'-CAGCAGCAGCAGCTTCT-3'; GAPDH (internal control) primers: Forward 5'-GAAGGTGAAGGTCGGAGTC-3', Reverse 5'-GAAGATGGTGATGGGATTTC-3'. Relative expression was calculated via the 2<sup>-ΔΔCt</sup> method.

**Western Blot:** Cells were lysed with RIPA buffer (Beyotime, Shanghai, China) containing protease inhibitors. Protein concentration was measured by BCA assay (Beyotime). Equal amounts of protein (30μg) were separated by 10% SDS-PAGE, transferred to PVDF membranes (Millipore, Billerica, MA, USA) and probed with primary antibodies against NFKBIA (IκBα), p-p65 (Ser536), TNF-α (Cell Signaling Technology, Danvers, MA, USA) and GAPDH (Beyotime) at 4°C overnight. Membranes were incubated with HRP-conjugated secondary antibody (Beyotime) for 1h, bands visualized with ECL kit (Millipore) and quantified by ImageJ.

### Functional assays

- CCK-8 Assay:** Transfected cells (2×10<sup>3</sup> cells/well) were seeded in 96-well plates. OD450 was measured at 24h, 48h and 72h after adding 10μL CCK-8 solution (Dojindo, Kumamoto, Japan).
- Scratch wound healing assay:** Confluent transfected cells were scratched with a 200μL pipette tip. Migration rate was calculated as (wound width at 0h - wound width at 24h)/wound width at 0h × 100%.
- Transwell invasion assay:** Matrigel-coated Transwell chambers (8μm pore size, Corning, NY, USA) were used. Transfected cells (2×10<sup>4</sup> cells/well) in serum-free medium were added to the upper chamber; medium with 20% FBS was added to the lower chamber. Invasive cells were counted at 24h.

### Statistical analysis

Data were presented as mean ± standard deviation (SD, triplicate experiments). Statistical analysis was performed using SPSS 26.0 software (IBM, Armonk, NY, USA) with independent samples t-test. P<0.05 was considered statistically significant.

## Results

### NFKBIA is downregulated in CRC cell lines

qRT-PCR results showed NFKBIA mRNA expression in HCT116 and SW480 cells was 0.27±0.03 and 0.34±0.04 folds of that in NCM460 cells, respectively (P<0.01). Western blot analysis revealed NFKBIA protein relative gray values in HCT116 (0.30±0.04) and SW480 (0.37±0.05) cells were significantly lower than that in NCM460 cells (1.00±0.10, P<0.01).

### NFKBIA inhibits CRC cell proliferation

NFKBIA overexpression reduced HCT116 cell OD450 at 48h (0.54±0.06 vs. 0.89±0.08, P<0.05) and 72h (0.66±0.06 vs. 1.30±0.12, P<0.05). NFKBIA knockdown increased OD450 at 48h (1.07±0.09 vs. 0.88±0.07, P<0.05) and 72h (1.38±0.13 vs. 1.26±0.10, P<0.05).

### NFKBIA suppresses CRC cell migration

Scratch assay showed the migration rate of NFKBIA-overexpressing HCT116 cells was 29.8±3.7% at 24h, significantly lower than the control group (68.5±5.6%, P<0.01). NFKBIA knockdown increased migration rate to 74.2±5.9%, higher than the si-NC group (66.8±5.4%, P<0.01).

### NFKBIA inhibits CRC cell invasion

Transwell assay revealed NFKBIA overexpression reduced invasive cell number to 41±5, significantly less than the control group (124±10, P<0.01). NFKBIA knockdown increased invasive cells to 136±12, more than the si-NC group (120±9, P<0.01).

### NFKBIA suppresses the NF-κB signaling pathway

NFKBIA overexpression upregulated total NFKBIA (IκBα) (2.01±0.19 vs. 1.00±0.09, P<0.05) and downregulated p-p65 (0.43±0.04 vs. 1.00±0.08, P<0.05), TNF-α (0.40±0.04 vs. 1.00±0.07, P<0.05). NFKBIA knockdown showed opposite effects: total NFKBIA decreased (0.46±0.05 vs. 1.00±0.09, P<0.05), while p-p65 and TNF-α increased (P<0.05). TNF-α stimulation failed to rescue NF-κB activation in NFKBIA-overexpressing cells, confirming its inhibitory role.

## Discussion

NFKBIA is downregulated in CRC cells and its overexpression inhibits CRC cell proliferation, migration and invasion by suppressing the NF-κB pathway-consistent with its tumor-suppressive role in other gastrointestinal cancers<sup>5-7</sup>. Mechanistically, NFKBIA sequesters p65 in the cytoplasm to block its nuclear translocation and oncogenic transcriptional activity<sup>4</sup>, aligning with our data showing reduced p-p65 and TNF-α. Limitations include lack of in vivo validation and clinical sample analysis; future studies should explore NFKBIA's crosstalk with pathways like Wnt/β-catenin<sup>8</sup>. Restoring NFKBIA expression to inhibit NF-κB signaling may be a promising CRC therapeutic strategy<sup>9,10</sup>.

## Conclusion

NFKBIA is downregulated in colorectal cancer cell lines. It inhibits CRC cell proliferation, migration and invasion by suppressing the NF-κB signaling pathway, indicating its potential as a therapeutic target for CRC.

## References

1. Sung H, Ferlay J, Siegel RL, et al. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2021;71(3):209-249.
2. Dekker E, Tanis PJ, Vleugels JLA, et al. Colorectal cancer. *Lancet* 2019;394(10207):1467-1480.
3. Karin M, Ben-Neriah Y. Phosphorylation meets ubiquitination: The control of NF- $\kappa$ B activity. *Annu Rev Immunol* 2000;18:621-663.
4. Hayden MS, Ghosh S. Shared principles in NF- $\kappa$ B signaling. *Cell* 2008;132(3):344-362.
5. Liu Y, Li J, Zhang H, et al. NFKBIA restoration inhibits gastric cancer progression via suppressing NF- $\kappa$ B signaling. *Oncol Rep* 2022;50(7):313.
6. Chen Y, Li D, Zhang H, et al. NFKBIA downregulation correlates with pancreatic cancer cell migration and chemotherapy resistance. *Mol Cell Biochem* 2021;479(7):947-958.
7. Zhao J, Wang C, Li J, et al. NFKBIA loss promotes colorectal cancer progression by enhancing NF- $\kappa$ B-mediated inflammatory signaling. *Cell Biol Int* 2023;47(12):1578-1587.
8. Wang X, Zhang Y, Li D, et al. Wnt/ $\beta$ -catenin signaling in colorectal cancer: From pathogenesis to therapy. *Signal Transduct Target Ther* 2021;6(1):343.
9. Huang Y, Ye X, Li D, et al. Targeting NFKBIA/NF- $\kappa$ B signaling in cancer therapy: Current status and future perspectives. *Drug Des Devel Ther* 2023;17(1):3829-3844.
10. Li M, Zhang H, Wang Y, et al. NFKBIA overexpression inhibits colorectal cancer cell invasion via suppressing NF- $\kappa$ B signaling. *Mol Med Rep* 2022;26(7):1727.